

EDITOR'S NOTE: The new, revised personal history and record card of physical findings is reproduced here for informational purposes. This is a card with the personal history on one side and the general health record on the opposite side. This card is the same size as the anesthesia record card. Both these record cards may be obtained by writing the office of the secretary, Dr. Richard I. Weaver, 316 Michigan Street, Toledo 2, Ohio. The price to members is \$3.00 for 500 cards.

PERSONAL HISTORY

Name Date

Address Record No.

Telephone Occupation

Age Sex Marital Status Name of Husband or Wife

Name of Family Physician Address

Telephone

Reason for Dental Visit.....

Date of Last Dental Visit Type of Treatment
(Reason)

Blood Pressure Pulse Urine Sugar Analysis

General Observations

.....

.....

.....

.....

.....

.....

GENERAL HEALTH RECORD

1. Are you, or have you been, under the care of a physician?
For what reason?
2. Have you ever been treated in a hospital? When?
For what condition?
3. Are you taking or have you ever taken any medicine or drugs?
Which one(s)? Have you ever
had a local anesthetic (novacain)? Did you ever have an unpleasant reaction
from one of them? Describe what happened
4. Have you ever had: (a) heart trouble; (b) rheumatic fever
(c) difficulty in breathing; (d) need for more than one pillow while sleeping
.....; (e) swollen ankles; (f) diabetes or thyroid trouble;
(g) epilepsy; (h) a severe digestive disturbance; (i) jaundice or
hepatitis; (j) an allergy (type); (k) hay fever or asthma
.....; (l) tuberculosis or syphilis; (m) anemia; (n) arthritis
.....; (o) any bleeding problems (type)
5. Do you smoke?; How much?

**Advance Registration 13th Annual Meeting
American Dental Society of Anesthesiology, Inc.
Essex Inn, Chicago, Illinois**

Dr. _____
(please print clearly)

Address _____

City _____ State _____ Zip _____

**THE SCIENTIFIC SESSION WILL COVER VARIOUS ASPECTS
OF PAIN CONTROL**

Program Chairman: DR. MILTON JAFFE

Newer General Anesthetic Agents Dr. Max Sadove
New Local Anesthetic Agents Dr. Harry Sicher
Role of Inhalation Analgesia Dr. Alvin Solomon
Role of Intravenous Agents Dr. Harold Marantz

Local Arrangements: DR. ELAINE STUEBNER

Dear Doctor Stuebner:

Please pre-register me for the National Meeting of American Dental Society of Anesthesiology. I will need hotel reservations at the Essex Inn, Chicago, for _____ starting:

- Feb. 25 } Business Meetings
- Feb. 26 }
- Feb. 27 Scientific Sessions (9 a.m. - 5 p.m., with luncheon)

- Member — no fee
- Non-Member — \$10.00
- Luncheon — \$6.00

(Tear out, check appropriate box and enclose total fee)

Scientific Session to be held at Conrad Hilton Hotel, across the street.



NEW ADSA MEMBERS

Dr. Didier Ardoin, II
University Hospitals
Cleveland, Ohio

Dr. Stanley D. Buckner
7001 North Clark Street
Chicago, Illinois 60626

Dr. Allen H. Cashman
1700 Grand Concourse
Bronx 57, New York

Dr. Donald M. Corbin
96-02 58th Avenue
Rego Park 68, New York

Dr. Douglas A. Drake
107 West First Street
Belvidere, Illinois

Dr. Sam Dyer
521 Madison
Maywood, Illinois

Dr. Manuel J. Freeman
2661 Salem Avenue
Dayton, Ohio

Dr. Irving M. Friedman
5616 Highland Road
Pontiac, Michigan

Dr. Rupert Q. Bliss
2720 Park Street
Jacksonville, Florida

Dr. William D. Bumsted
2811 L Street
Sacramento, California

Dr. Bert F. Erwin
Cape Royal, Suite 905
Cocoa Beach, Florida

Dr. James G. Ghostley
603 Bemidji Avenue
Bemidji, Minnesota


Dr. Franklyn A. Goodell
516½ Bettrami Avenue
Bemidji, Minnesota

Dr. Charles L. Harer
Nekoosa Medical Center
Nekoosa, Wisconsin

100
'EMPIRIN'®
COMPOUND
FAST RELIEF OF HEADACHE, TOOTHACHE, MUSCULAR
ACHES AND PAINS, DISCOMFORT OF COLDS.
Each tablet contains phenacetin 2½ gr., as-
pirin 3½ gr., caffeine ½ gr.
DOSE: Adults, 1 or 2 tablets with water. Re-
peat every 3 hours as needed. Do not exceed
6 tablets per day. Children 4-12 years, ½
adult dose; under 4, consult your doctor.
WARNING: Do not exceed recommended dose,
or take regularly for longer than 10 days un-
less directed by a doctor. If used in large
amounts and for a long period of time, may
be harmful to the kidneys. Keep this and all medicines out of children's
reach.
BURROUGHS WELLCOME & CO.
(U.S.A.) Inc., Tuckahoe, N.Y.
1240 Made in U.S.A.

**A good analgesic
for your patients
who need
dependable
pain control
—relieves pain
from toothache
or tooth
extraction**

Also available: 'Empirin'® Compound with Codeine Phosphate* No. 1, gr. ¼ — No. 2, gr. ¼ — No. 3, gr. ½
— No. 4, gr. 1 *Warning may be habit forming. Available on oral R_x where state law permits.



16mm.
scientific films
available
on a loan basis.

These films are available without charge to dentists, hospitals, professional societies and teaching institutions. Reservations should be made at least three weeks in advance, and should include an alternate film title. Address your request to the Astra Film Library, giving the name of your organization, type of audience, and the name, position and address of the person to receive the films. All postage will be prepaid.

1. Regional Block Anesthesia in Dentistry by M.D. Anderson Hospital, University of Texas, Houston, Texas. Color, sound, 19 minutes. Appropriate anatomy and clinical demonstrations of various dental nerve blocks are shown including: Posterior, middle, and anterior superior alveolar, posterior palatine, nasopalatine, infra-orbital, inferior alveolar, lingual, long buccal, and mental.

2. Minimum Dosage Local Anesthesia by Adrian Cowan, M.B., F.D.S.R.C.S., England. Color, sound, 45 minutes. Anatomical landmarks for infiltration and nerve block are demonstrated clinically. Onset times, extent, duration and volumes used are discussed for each procedure and an apicectomy, extraction and gingivectomy are shown.

3. Frenectomy by Lester M. Silverman, B.A., D.D.S., M.S.D. Black and white kinescope, sound, 26 minutes. Two clinical cases with abnormal labial frenums are used to demonstrate the frenectomy technique for both orthodontic and prosthetic purposes.

4. Regional Analgesia for Surgery of the Head and Neck by William S. Derrick, M.D. Color, sound, 27 minutes. An anatomical study of the head and neck followed by a demonstration of the following nerve blocks: Superficial and deep cervical, mandibular, maxillary, mental, infra-orbital, and medial orbital.

5. Transtracheal Anesthesia by Charles C. Wycoff, M.D. Color, sound, 24 minutes. Anatomical landmarks are exposed by dissection of the pharyngolaryngeal region. Patients are injected, scoped, and intubated. Nasal endotracheal intubation and superior laryngeal injection of the patients are demonstrated.

6. Case Management, Boston University School of Medicine, Boston, Massachusetts.

Part I. Examination by Henry M. Goldman, D.M.D., F.A.C.D., F.I.C.D. and Gerald Kramer, D.M.D., F.A.C.D. Color, sound, 18 minutes. Examination of a dentition affected by loss of teeth, caries, a periapical lesion at the apex of a maxillary right lateral incisor, and periodontal disease.

Part II. Initial Preparation in Periodontics by Henry M. Goldman, D.M.D.,

F.A.C.D., F.I.C.D. and Gerald Kramer, D.M.D., F.A.C.D. Color, sound, 33 minutes. Demonstration of two essential phases of initial therapy—scaling and the teaching of oral physiotherapy.

Part III. The Surgical Phase of Periodontal Therapy by Henry M. Goldman, D.M.D., F.A.C.D., F.I.C.D. and Gerald Kramer, D.M.D., F.A.C.D. Color, sound, 43 minutes.

Demonstrates the technique of gingivectomy as part of periodontal therapy. The preoperative status, operative procedure and postoperative technique of packing and related problems are shown.

Part IV. Endodontics—Non-Surgical Treatment by Herbert Schilder, D.D.S. Color, sound, 48 minutes. Demonstrates non-surgical treatment of a maxillary lateral incisor which has undergone pulpal degeneration, and includes cleaning, shaping, sterilization and packing of the root canal.

Part V. Clinical Procedures for Fixed Bridge Prosthesis by David J. Baraban, D.M.D., F.A.C.D. Color, sound, 47 minutes. Tooth preparation, temporary and final restorations, including impressions in fixed bridge prosthetics.

Astra Pharmaceutical Products, Inc.
Worcester, Massachusetts 01606

ASTRA®
©1965 ASTRA



Lidocaine,
the local
anesthetic
you trust!

Now,
in the
sterile system
you require!

IT'S NEW!

Graham
LIDOCAINE
DISPOSALL®

STERILE NEEDLE & CARTRIDGE
"ALL-IN-ONE" SEALED UNIT



Now, Graham makes LIDOCAINE available in its DISPOSALL® cartridges — the only effective closed system ever designed.

YOU GET ASSURED ASEPSIS WITH THE RESULTS YOU'VE COME TO EXPECT.

Simply insert the DISPOSALL® into the syringe, remove the needle guard and administer. Chances for cross infection are virtually eliminated.

Graham's LIDOCAINE DISPOSALLS® with epinephrine 1:50,000 and 1:100,000 are available in 27 gauge long and short needles; 23 and 25 gauge long only. Introductory Offer Available.

LIDOCAINE also available in regular cartridges from your dental dealer.

Graham CHEMICAL CORP., Dept. E-12, 129-21 Merrick B'ld., Springfield Gardens, L.I., N.Y.



PENTHRANE® (methoxyflurane)

for easy maintenance with outstanding safety


Here is why Penthrane anesthesia permits such good safety in oral surgery and operative dentistry:

1. It is free of fire and explosion hazard.
2. Small alterations in Penthrane vapor concentration do not precipitate abrupt physiologic responses.
3. Excellent analgesia can be attained with a relatively light plane of anesthesia.
4. Blood pressure and heartbeat are remarkably stable.
5. Toxicity is low, and there is no interference with hemostasis.
6. Low doses suffice for maintenance, and also favor a rapid recovery.

Used in conjunction with Pentothal (thiopental) to speed induction, Penthrane is of great use to the dental anesthesiologist in all but the shorter procedures.

PRECAUTIONS Use only conservative doses of epinephrine, barbiturates, narcotics, tubocurarine, or trimethaphan camphorsulfonate. Suspected liver disease may rule out halogenated agents such as Penthrane. Guedel eye signs do not apply. Blood pressure, respiratory depth, and muscular relaxation are of most value in estimating depth of anesthesia.

502195



PENTOTHAL® (thiopental) for pleasant, rapid induction


Pentothal is an agent of choice for anesthetic induction. Note its many advantages for the dental anesthesiologist:

1. It is rapid-acting. Sleep comes within seconds.
2. Pentothal is pleasant to the patient. It provides amnesia, and does not produce a delirium stage.
3. Narcosis is easily regulated, and can be increased with great speed if needed.
4. Coughing, sneezing, or other troublesome respiratory irritation is seldom associated with it.
5. Its action is brief at the doses that suffice for induction, permitting prompt recovery.

Pentothal has proved ideal for use with Penthrane, where it brings ample speed of induction. In short procedures, Pentothal may also be used as a sole anesthetic. We invite you to use it and Penthrane in your own practice.

PRECAUTIONS Pentothal is contraindicated in severe respiratory embarrassment, liver or kidney disease, advanced cardiac disease, and prophyria. Use it with caution in the presence of other disabilities.

502197



Presenting the new needleless and painless


ROBBINS DENTJECTOR *

The only authentic European automatic dental pressure injection syringe for oral anesthetic solutions.

The application of this time-tested, unique, anesthetic apparatus makes it possible, for the first time, to achieve an instantaneous and painfree nerve blockage, in a wide range of dental-surgical modalities without the hypodermic needle.

The DENTJECTOR technique achieves *painless* tissue infiltration with a high-velocity microspray, using quick-blocking anesthetic solutions in single or multiple dose of .1 cc. to a depth of 2 - 5 mm. *without* contact with the site of injection.

The DENTJECTOR accepts any sterilization method up to 284° F., and is guaranteed for one year from the day of purchase.



What dentists are reporting about the DENTJECTOR:

"Injections are painless and have received very good patient acceptance."

THOMAS C. MATTHEWS, D.D.S., Oregon, Ohio.

"Everyone is enthusiastic about it."

CLIFFORD E. FISHER, D.D.S., Lubbock, Texas.

"A remarkable instrument."

DAVID LASSER, D.D.S., Brooklyn, New York.


ROBBINS INSTRUMENTS COMPANY

EXCLUSIVE NATIONAL DISTRIBUTORS
DIRECT SALES AND SERVICE

CHATHAM, NEW JERSEY 07928

*U.S. Patent No. 3,129,708

The only head with all 7 controls



McKesson Seeler Resuscitator Head features complete fingertip control

This versatile head can be used with any mask or endotracheal tube without adapters. Connected to an at-hand or remote oxygen supply, you always have complete control at your fingertips.

1. Gas flow valve adjusts from off to maximum. Cycles per minute increase with flow increase.

2. Oxygen diluter valve adjusts from 50 to 100% oxygen concentrations. Important with portable resuscitators.

3. Negative pressure control can be adjusted from 0 to -7mm of Hg.

5. Resuscitator valve turns counter clockwise so unit can be used as an inhalator.

6. Positive pressure control can be adjusted from $+7\text{mm}$ to $+15\text{mm}$ of Hg.

4. & 7. Manual overrides for positive and negative pressure enables operator to ventilate lungs when airway passages are blocked or closed - chest cardiac massage is performed.

WRITE TODAY FOR COMPLETE INFORMATION.

McKESSON APPLIANCE CO.

1500 W. VAN DE GRAFF ST.
ANN ARBOR, MICH. 48106



McKesson
Appliance Co.
Ann Arbor, Mich.



Decisions, decisions

With the two formulations of Carbocaine you have a choice—a choice that actually minimizes decision-making...since the case at hand usually makes the decision for you.

All cases, of course, require anesthesia which is rapid in onset, profound in effect and well tolerated.

But what about duration? Would relatively short operating anesthesia be desirable?

Decision: Carbocaine 3% without vasoconstrictor.

Is the procedure complex, or does it present other facets which dictate extended duration?

Decision: Carbocaine 2% with Neo-Cobefrin 1:20,000.

The fact that Carbocaine, unlike any other local anesthetic for dental purposes¹, needs a vasoconstrictor only to extend duration and not to potentiate depth, gives you a choice...a simple decision to make.

FOR YOUR INFORMATION: These local anesthetic solutions are for dental block and infiltration injections only. There are no known contraindications in dentistry except for patients known to be sensitized. Inject slowly and avoid intravascular injection by aspirating. As with all local anesthetic solutions, adverse reactions due to intolerance, overdosage, or intravascular injection may occur and include hypotension and respiratory depression. Such reactions occur infrequently and usually are readily controlled by supporting circulation with a vasopressor and respiration with oxygen.

Carbocaine and Neo-Cobefrin are trademarks (Reg. U.S. Pat. Off.) of Sterling Drug Inc.

Carbocaine[®] HCl

brand of mepivacaine HCl

Carbocaine HCl 2% with Neo-Cobefrin[®] 1:20,000
(brand of levo-nordefrin)

rapid-acting, consistently profound anesthesia of sufficiently long duration that even extensive and complicated procedures seldom require reinjection.

Carbocaine HCl 3% without vasoconstrictor provides very rapid, profound anesthesia without long duration—particularly desirable in children's dentistry and modern high-speed restorative procedures. *Operating anesthesia averages 20 minutes upper, 40 minutes lower.*

Reference: 1. Weil, C., Santangelo, C., Welham, F. S., and Yackel, R. F. Clinical evaluation of mepivacaine hydrochloride by a new method. J.A.D.A. 63:26, July 1961.

Cook-Waite Laboratories, Inc., New York, N.Y.

COOK WAITE